Valorization of Methyl Azelaaldehyde - A Vegetable Oil Based Platform Molecule for the Synthesis of Monomers through Stetter Reaction

Aubin Charvieux, Nam Duc Vu, Nicolas Duguet, Marc Lemaire

► To cite this version:

Aubin Charvieux, Nam Duc Vu, Nicolas Duguet, Marc Lemaire. Valorization of Methyl Azelaaldehyde - A Vegetable Oil Based Platform Molecule for the Synthesis of Monomers through Stetter Reaction. European Journal of Organic Chemistry, 2019, 2019 (6), pp.1251-1256. 10.1002/ejoc.201801611 . hal-02128677

HAL Id: hal-02128677

<https://hal-udl.archives-ouvertes.fr/hal-02128677>

Submitted on 3 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Valorization of methyl azelaaldehyde – a vegetable oil-based platform molecule – to monomers through Stetter reaction

Aubin Charvieux,^[a] Nam Duc Vu,^[a] Nicolas Duguet^{*[a]} and Marc Lemaire^{*[a]}

Abstract: The valorization of vegetable oil-derived methyl azelaaldehyde (methyl 9-oxo-nonanoate) to monomers was studied through NHC-catalysed Stetter reaction. Among the Michael acceptors tested, dimethyl fumarate gave the highest selectivity (97 %) for the corresponding Stetter adduct, thus limiting the competing benzoin condensation.

Introduction

Biomass has received considerable attention in the recent years as a sustainable alternative to petroleum derivatives.^[1] In this regard, vegetable oils represent a valuable feedstock as they are widely available and relatively inexpensive. Contrary to lignin or carbohydrates, they have low oxygen content, thus making them perfectly suitable for biofuel applications.^[2] However, these valuable resources can be best transformed to bio-based platform chemicals^[3] and polymers^[4] with higher added values. This strategy is also beneficial from an environmental point of view, as it incorporates CO₂ in materials instead of releasing it in the atmosphere.

Three main approaches are reported in the literature to convert vegetable oils to polymers: i) unsaturated vegetable oils and their corresponding epoxidized derivatives can be directly (co-)polymerized to give functional materials such as linoleum, alkyd resins or epoxy resins; ii) unsaturated fatty acids can be also functionalized to produce monomers through a wide range of transformations; iii) unsaturated fatty acids can be cleaved to give shorter fragments that can be further transformed to polymer building-blocks. This last strategy is particularly attractive as short monomers (C6-C12) usually give access to materials with better mechanical properties. For example, methyl ricinoleate obtained from castor oil can be thermally cracked to give heptanal and methyl 10-undecenoate that is further transformed to nylon-11 (Rilsan®).^[5] Terminal alkenes can also be produced by cross-metathesis of unsaturated fatty acid derivatives with ethylene (ethenolysis).^[6] The alkenes can be further transformed to the corresponding homologated aldehydes through hydroformylation.^[7] Advantageously, functionalized aldehydes can be directly obtained by reductive

ozonolysis of unsaturated fatty acid derivatives.^[8] Specifically, the reductive ozonolysis of methyl oleate gives nonanal and methyl azelaaldehyde (methyl 9-oxo-nonanoate) as cleavage compounds and each of them could be valorized following their own value chain. Nonanal can be used as such or serves as an alkylating agent for the preparation of 100% biobased surfactants through aldolisation^[9] or reductive alkylation (reductive etherification) of polyols^[10] and sugar derivatives.^[11] Comparatively, there are not so many ways to valorize methyl azelaaldehyde. On the one hand, the reduction of its aldehyde function gives the corresponding alcohol and the resulting monomer - methyl 9-hydroxynonanoate - can be further (co-)polymerized to give polyesters (Scheme 1, a).^[12] On the other hand, the reductive amination with ammonia affords methyl 9-aminononanoate that serves as a monomer for the preparation of polyamide-9 or mixed polyamides (Scheme 1, b).^[13] In this context, complementary routes to valorize methyl azelaaldehyde are of great interest.

Scheme 1. Valorization of methyl azelaaldehyde to monomers.

Recently, organocatalysis has emerged as a new tool for the valorization of biomass.^[14] Among available organocatalysts, *N*-Heterocyclic carbenes (NHCs)^[15] occupy a unique place as they can promote *Umpolung* reactivity.^[16] This specific reactivity has been notably exploited for the valorization of bio-based aldehydes such as furfural, 5-hydroxymethylfurfural (HMF)^[17] and bioethanol-derived acetaldehyde through Benzoin condensation.^[18] The Stetter reaction,^[19] i.e. the 1,4-addition of an aldehyde onto a Michael acceptor, also employs *Umpolung* reactivity and has been extensively studied in both intra-^[20] and intermolecular versions.^[21] However, it has been by far less studied in the context of the valorization of biomass. To the best of our knowledge, only Chi *et al.* have reported the use of the Stetter reaction to trap formaldehyde equivalents – generated from hexoses through retro-benzoin condensation – with enones to give the corresponding β -formylketones.^[22] In this context, we report herein a complementary route for the valorization of methyl azelaaldehyde by NHC-catalysed Stetter reaction with acrylates to give functionalized monomers (Scheme 1, c).

[a] A. Charvieux, N. D. Vu, Dr N. Duguet, Pr M. Lemaire
Univ Lyon, Université Claude Bernard Lyon1, CNRS, INSA, CPE-
Lyon, Institut de Chimie et Biochimie Moléculaires et
Supramoléculaires, ICBMS, UMR 5246, Equipe Catalyse, Synthèse
et ENvironnement (CASYEN), Bâtiment Lederer, 1 rue Victor
Grignard, F-69100 Villeurbanne cedex, France.
Phones: +33(0)-472-448-507, +33(0)-472-431-407.
E-mails: nicolas.duguet@univ-lyon1.fr, marc.lemaire.chimie@univ-lyon1.fr

Supporting information for this article is given via a link at the end of the document.

Results and Discussion

Methyl azelaaldehyde can be prepared from methyl oleate through reductive ozonolysis. However, this method requires specific equipment and is dangerous due to the toxicity of ozone and the formation of explosive intermediates. That is the reason why, it was synthesized following our previously reported procedure (Scheme 2).

Scheme 2. Preparation of methyl azelaaldehyde from methyl oleate.

First, methyl oleate **1** (96% purity) was epoxidized to methyl 9,10-epoxystearate **2** in 96% isolated yield.^[23] The epoxide was then hydrolyzed, giving methyl 9,10-dihydroxystearate **3** in 90% isolated yield. This diol was selectively oxidized using a Pd(OAc)₂ / neocuproine complex under oxygen atmosphere to give α -hydroxyketone **4** with 80% yield.^[24] Alternatively, it can also be prepared by dehydrogenation of diol **3** using Ru/C with removal of hydrogen under vacuum.^[24] Finally, the cleavage of **4** through NHC-catalysed retro-benzoin process gave aldehydes **5** and **6** under reactive distillation conditions, with 85 and 71% yields, respectively.^[25]

With methyl azelaaldehyde **6** (GC purity = 98%) in hands, we envisioned that the Stetter reaction using a range of acrylates as Michael acceptors would provide multi-functional compounds that could serve as monomers for polymer applications. To find appropriate reaction conditions, methyl acrylate was selected as a model substrate. The Stetter reaction between methyl azelaaldehyde **6** and methyl acrylate was already reported by Novák *et al.*^[26] The Stetter adduct was obtained in only 42% yield. However, the corresponding symmetrical α -hydroxyketone was also formed in 58% yield, showing that the benzoin condensation is a competing reaction. This reaction has not been further optimized by the authors as the desired product was only an intermediate in the preparation of prostaglandin synthons.

From a mechanistic point of view, the Stetter reaction and the benzoin condensation both occur through the common Breslow intermediate^[27] **III** formed by the nucleophilic addition of carbene **I** onto the aldehyde (Scheme 3). Thus, the selectivity between Stetter and benzoin products is directly correlated to the reaction between the Breslow intermediate and the present electrophiles (aldehyde or Michael acceptor). In most cases, the aldehyde is

by far more reactive than the Michael acceptor, leading to the α -hydroxyketone as primary product. However, considering that the benzoin condensation is a reversible process while the Stetter reaction is not, the Stetter adducts can be produced as final products, as these species are usually thermodynamically favored.^[28] With aromatic aldehydes ($R^1 = \text{Ar}$), the reverse process can occur at low temperature (typically 20°C).

Scheme 3. Stetter and benzoin condensation competing reaction mechanisms.

Consequently, aromatic α -hydroxyketones, *i.e.* benzoin, can be used as aldehyde precursors for the Stetter reaction.^[29] In contrast, the α -hydroxyketones produced from aliphatic aldehydes ($R^1 = \text{alkyl}$), *i.e.* acyloins, are by far more stable than their aromatic analogues. Therefore, the retro-acyloin process can only occur at elevated temperature (typically 130°C) and can only be catalyzed by thiazolide species generated from the corresponding thiazolium salts, as reported by Miyashita,^[30] Chi^[22] and our group,^[25] and supported by DFT calculations.^[31]

From these previous results and mechanistic considerations, we have first investigated the Stetter reaction of methyl azelaaldehyde **6** with methyl acrylate using a small range of thiazolium precatalysts at 130°C under microwave irradiation (Table 1).^[32]

With commercially available thiamine-derived thiazolium salt **10** (20 mol%) and K₂CO₃ (10 mol%), the conversion of **6** reached 59% and Stetter adduct **8** was obtained with only 27% yield (Table 1, entry 1). Under these conditions, α -hydroxyketone **9** was also formed with 18% yield. With our home-made^[25b] thiazolium triflate **11**, the conversion improved to 89% and **8** was obtained with 48% yield (Table 1, entry 2). Slightly better results were obtained with commercially-available thiazolium **12**, giving **8** with 50% yield (Table 1, entry 3). Finally, triazolium salt **13**,

used as negative control, gave no Stetter product and only yielded the self-aldolisation of aldehyde **6**, confirming again the superiority of thiazolylidene species to catalyze the desired transformation (Table 1, entry 4). From these results, thiazolium salt **12** was chosen as a precatalyst for further optimization.

Table 1. Azolium precatalyst screening.^[a]

6 (1 equiv.) + 7 (10 equiv.)

azolium salts **10-13** (20 mol%)
 base (n mol%)
 dry CH₃CN
 130°C (μW), 30 min

8 + 9

Entry	Azolium salt	Conv. ^[b] [%]	Yield ^[b] [%]	
			6	9
1		59	27	18
2		89	48	28
3		92	50	33
4		56	0	0

[a] Conditions: microwave tube, aldehyde **6** (1 equiv., 1 mmol), **7** (10 equiv.), azolium salt (20 mol%), K₂CO₃ (10 mol%) for precatalyst **10-12** or tBuOK (20 mol%) for precatalyst **13**, dry CH₃CN (1.5 mL), 130°C (μW), 30 min; [b] the conversions and yields were determined by GC using hexadecane as internal standard.

The effect of the quantity of Michael acceptor **7** has then been next studied (Table 2). As expected, the yield in Stetter product **8** progressively increased between 2.6 and 10 equivalents of methyl acrylate (Table 2, entries 1-3). Further increase to 15 equivalents did not lead to significant improvement, as the selectivity only increased slightly (Table 2, entry 4). Consequently, 10 equivalents of methyl acrylate were selected as a good compromise. Finally, increasing the reaction time under our best conditions (Table 2, entry 3) did not improve significantly the results. Overall, under the best conditions, the desired Stetter product **8** was afforded with only 50% yield, while the undesired acyloin **9** was obtained in 33% yield. This moderate selectivity could be explained by the low electrophilicity of methyl acrylate. This assumption is supported by Domingo *et al.* in a DFT study aiming at classifying several Michael acceptors according to their reactivity toward nucleophilic addition.^[33] Keeping in mind that the objective is to synthesize aliphatic Stetter products with suitable functional

groups for polymerization, a range of Michael acceptors was tested under the best conditions (Table 3). Fumaric acid **14** did neither give the Stetter product nor the acyloin (Table 3, entry 1). This can be explained by its insolubility in acetonitrile and the fact that it can react with the base, thus preventing the formation of the active carbene species.

Table 2. Effect of the quantity of Michael acceptor.

Entry	Quantity of 7 [equiv.]	Conv. ^[b] [%]	Yield ^[b] [%]	
			8	9
1	2.6	93	23	52
2	5	91	37	43
3	10	92	50	33
4	15	83	52	26

[a] Conditions: microwave tube, aldehyde **6** (1 equiv., 1 mmol), **7**, thiazolium salt **12** (20 mol%), K₂CO₃ (10 mol%), dry CH₃CN (1.5 mL), 130°C (μW), 30 min; [b] the conversions and yields were determined by GC chromatography using hexadecane as internal standard.

Surprisingly, no reaction occurred with maleic anhydride **15**, though this compound is one of the most electrophilic Michael acceptors (Table 3, entry 2).^[33,34] Consequently, we hypothesized that the carbene formed from thiazolium **12** could directly add onto maleic anhydride instead of the aldehyde, thus giving a stable species and trapping the catalyst. Fumaronitrile **16** gave a high selectivity for the Stetter adduct **22** (94%) but this result is not very representative considering that the conversion was very low (Table 3, entry 3). With dimethyl itaconate **17**, the conversion improved to 27% but almost no selectivity was observed (Table 3, entry 4). The use of dimethyl maleate **18** and acrylonitrile **19** gave good conversions (88%) but moderate selectivity for the Stetter products **24** and **25** (50 and 72% respectively) (Table 3, entries 5-6). Slightly better results were obtained with methyl acrylate under these reaction conditions (Table 3, entry 7). Dimethyl fumarate **20** gave a good 85:15 selectivity for the triester **24** but the conversion remained modest (65%) (Table 3, entry 8). Satisfyingly, when using 5 equivalents of **20**, the selectivity towards the Stetter adduct **24** improved to 97% after 1 hour (Table 3, entry 9). Moreover, when using chalcone **21** as a starting material, the corresponding Stetter adduct **26** was obtained with 93% selectivity at complete conversion (Table 3, entry 10). These results show that the selectivity could be switched in favor of the Stetter adduct, even with aliphatic aldehydes, provided a suitable Michael acceptor. In our case, the most suitable Michael acceptors should have an

Scheme 5. Proposed mechanism for the formation of deoxy-Breslow intermediate.

To test this hypothesis, the reaction was conducted under similar conditions but without aldehyde **6**. Under these conditions, diester **28** was observed in the crude ^1H NMR spectrum, thus confirming our proposal. Interestingly, diester **28** could be attributed to a deoxy-Breslow intermediate. Deoxy-Breslow intermediates of triazolylidene and imidazolylidene species have been already reported when using alkyl halides,^[36] acrylates^[37] or chalcones.^[38] However, to the best of our knowledge, deoxy-Breslow species have never been observed with thiazolylidene species. Consequently, the catalytic activity of **28** has been probed on the Stetter reaction under the same conditions reported in Table 3, entry 8 (with or without K_2CO_3). However, no Stetter adduct nor benzoin product was obtained under these conditions, indicating that **28** is relatively stable under our reaction conditions (130°C) and only zwitterionic intermediate **27** could reverse to release the free carbene.

Conclusions

In conclusion, we have developed a new valorization route of vegetable oil-derived methyl azelaaldehyde (methyl 9-oxononanoate) through NHC-catalysed Stetter reaction. It was found that dimethyl fumarate gave the best results with an excellent conversion (95%) and a high selectivity (97%) for the Stetter product, thus limiting the competing benzoin condensation. The di- and triesters obtained through this methodology could serve as building-blocks for the preparation of bio-based polymers.

Experimental Section

General procedure for Stetter reaction under microwave activation:

In a 5 mL microwave tube, methyl azelaaldehyde (**6**; 200 mg, 1.07 mmol, 1 equiv.), NHC precursor (**10-13**; 0.215 mmol, 0.20 equiv.), K_2CO_3 (15 mg, 0.11 mmol, 0.10 equiv.), Michael acceptor in desired quantity (**7,14-21**) were combined. Dry CH_3CN (1.5 or 2 mL) was added. The mixture was stirred under microwave irradiation at desired temperature, for a period of time. The mixture was cooled to room temperature and analysed by GC (with or without hexadecane as an internal standard).

Acknowledgements

The authors thank the SAS PIVERT for a Ph.D. grant to N. D. Vu and a master grant for A. Charvieux. (GENESYS program – project WP3P21-Bioaldehydes). This work was performed in partnership with the SAS PIVERT, within the frame of the French Institute for the Energy Transition (Institut pour la Transition Énergétique (ITE) P. I. V. E. R. T. (<http://www.institut-pivert.com>)) selected as an Investment for the Future ("Investissements d'Avenir"). This work was supported, as part of the Investments for the Future, by the French Government under the reference ANR-001-01.

Keywords: Aldehydes • Monomers • Stetter reaction • Umpolung • Vegetable oils

- [1] a) P. Gallezot, *Chem. Soc. Rev.*, **2012**, *41*, 1538–1558. b) A. Corma, S. Iborra and A. Velty, *Chem. Rev.*, **2007**, *107*, 2411–2502. c) C. O. Tuck, E. Pérez, I. T. Horváth, R. A. Sheldon and M. Poliakoff, *Science*, **2012**, *337*, 695–699.
- [2] a) G. W. Huber, S. Iborra and A. Corma, *Chem. Rev.*, **2006**, *106*, 4044–4098. b) M. Mittelbach, *Eur. J. Lipid Sci. Technol.*, **2015**, *117*, 1832–1846. c) T. Issariyakul and A. K. Dalai, *Renewable Sustainable Energy Rev.*, **2014**, *31*, 446–471.
- [3] U. Biermann, U. Bornscheuer, M. A. R. Meier, J. O. Metzger and H. J. Schäfer, *Angew. Chem. Int. Ed.*, **2011**, *50*, 3854–3871.
- [4] a) G. Lligadas, J. C. Ronda, M. Galià and V. Cádiz, *Materials Today*, **2013**, *16*, 337–343. b) A. Gandini, T. M. Lacerda, A. J. F. Carvalho and E. Trovatti, *Chem. Rev.*, **2016**, *116*, 1637–1669.
- [5] N. D. Spiccia, E. Border, J. Illesinghe, W. R. Jackson and A. J. Robinson, *Synthesis* **2013**, *45*, 1683–1688.
- [6] G. S. Forman, R. M. Bellabarb, R. P. Tooze, A. M. Z. Slawin, R. Karch, R. Winde, *J. Organomet. Chem.* **2006**, *691*, 5513–5516; b) K. A. Burdett, L. D. Harris, P. Margl, B. R. Maughon, T. Mokhtar-Zadeh, P. C. Saucier, E. P. Wasserman, *Organometallics* **2004**, *23*, 2027–2047; c) R. M. Thomas, B. K. Keitz, T. M. Champagne, R. H. Grubbs, *J. Am. Chem. Soc.* **2011**, *133*, 7490–7496; d) J. C. Mol, *Green Chem.* **2002**, *4*, 5–13.
- [7] a) A. Behr, A. J. Vorholt. Hydroformylation and Related Reactions of Renewable Resources. In: Meier M., Weckhuysen B., Bruijninx P. (eds) *Organometallics and Renewables. Top. Organomet. Chem.*, **2012**, *39*, 103–128. Springer, Berlin, Heidelberg; b) M. Furst, V. Korkmaz, T. Gaide, T. Seidensticker, A. Behr, A. J. Vorholt, *ChemCatChem*, **2017**, *9*, 4319–4323.
- [8] a) K. Louis, L. Vivier, J.-M. Clacens, M. Brandhorst, J.-L. Dubois, K. De Oliveira Vigier, Y. Pouilloux, *Green Chem.* **2014**, *16*, 96–101; b) T. S. Omonov, E. Kharraz, P. Foley, J. M. Curtis, *RSC Adv.* **2014**, *4*, 53617–53627; c) M. D. Lundin, A. M. Danby, G. R. Aken, T. P. Binder, D. H. Busch, B. Subramaniam, *ACS Sustainable Chem. Eng.* **2015**, *3*, 3307–3314.
- [9] a) B. Zhu, D. Belmessieri, J. F. Ontiveros, J.-M. Aubry, G.-R. Chen, N. Duguet, M. Lemaire, *ACS Sustainable Chem. Eng.*, **2018**, *6*, 2630–2640; b) B. Zhu, G.-R. Chen, N. Duguet, M. Lemaire, *ACS Sustainable Chem. Eng.* **2018**, *6*, 11695–11703.
- [10] Y. Shi, W. Dayoub, A. Favre-Régouillon, G.-R. Chen, M. Lemaire, *Tetrahedron Lett.*, **2009**, *50*, 6891–6893; b) Y. Shi, W. Dayoub, G.-R. Chen, M. Lemaire, *Green Chem.* **2010**, *12*, 2189–2195; c) Sutter, M.; Da Silva, E.; Duguet, N.; Raoul, Y.; Métay, E.; Lemaire, M. *Chem. Rev.* **2015**, *115*, 8609–8651; d) A. Herbinski, E. Métay, E. Da Silva, E. Illous J.-M. Aubry, M. Lemaire, *Green Chem.*, **2018**, *20*, 1250–1261.
- [11] a) C. Gozlan, R. Lafon, N. Duguet, A. Redl, M. Lemaire, *RSC Adv.* **2014**, *4*, 50653–50661; b) Gozlan, C.; Deruer, E.; Duclos, M.-C.; Molinier, V.; Aubry, J.-M.; Redl, A.; Duguet, N.; Lemaire, M. *Green Chem.* **2016**, *18*, 1994–2004; c) Belmessieri, D.; Gozlan, C.; Duclos, M.-C.; Molinier, V.; Aubry, J.-M.; Dumitrescu, O.; Lina, G.; Redl, A.; Duguet, N.; Lemaire, M. *Eur. J. Med. Chem.* **2017**, *128*, 98–106; d) Belmessieri,

- D.; Gozlan, C.; Duclos, M.-C.; Dumitrescu, O.; Lina, G.; Redl, A.; Duguet, N.; Lemaire, M. *Bioorg. Med. Chem. Lett.* **2017**, *27*, 4660–4663.
- [12] a) Z. S. Petrović, J. Milić, Y. Xu, I. Cvetković, *Macromolecules*, **2010**, *43*, 4120–4125; b) J. Jose, G. Pourfallah, D. Merkley, S. Li, L. Bouzidi, A. Lopes Leao and S. S. Narine, *Polym. Chem.*, **2014**, *5*, 3203–3213; c) J. Jose, G. Pourfallah, A. Lopes Leao and S. S. Narine, *Polym. Int.*, **2014**, *63*, 1902–1911.
- [13] W. L. Kohlhasse, E. H. Pryde, J. C. Cowan, *J. Am. Oil Chem. Soc.*, **1970**, *47*, 183–188.
- [14] For reviews on the organocatalytic transformations of biomass and CO₂, see: a) D. Liu, E. Y.-X. Chen, *Green Chem.* **2014**, *16*, 964–981; b) G. Fiorani, W. Guo, A. W. Kleij, *Green Chem.* **2015**, *17*, 1375–1389.
- [15] For reviews on NHC catalysis including benzoin condensation, see: a) X. Bugaut, F. Glorius, *Chem. Soc. Rev.* **2012**, *41*, 3511–3522; b) C. D. Campbell, K. B. Ling, A. D. Smith, in *N-Heterocyclic Carbenes in Transition Metal Catalysis and Organocatalysis* (Ed. C. S. J. Cazin), Springer, Netherlands, **2011**, pp. 263–297; c) P.-C. Chiang, J. W. Bode, in *N-Heterocyclic Carbenes: From Laboratory Curiosities to Efficient Synthetic Tools* (Ed. S. Díez-González), Royal Society of Chemistry, Cambridge, **2010**, pp. 339–445; d) E. M. Phillips, A. Chan, K. A. Scheidt, *Aldrichimica Acta* **2009**, *42*, 55–66; e) D. Enders, O. Niemeier, A. Henseler, *Chem. Rev.* **2007**, *107*, 5606–5655; f) N. Marion, S. Díez-González, S. P. Nolan, *Angew. Chem. Int. Ed.* **2007**, *46*, 2988–3000.
- [16] D. Seebach and D. Enders, *Angew. Chem., Int. Ed. Engl.*, **1975**, *14*, 15–32.
- [17] a) D. Liu, Y. Zhang, E. Y.-X. Chen, *Green Chem.* **2012**, *14*, 2738–2746; b) D. Liu, E. Y.-X. Chen, *ChemSusChem* **2013**, *6*, 2236–2239; c) B. L. Wegenhart, L. Yang, S. C. Kwan, R. Harris, H. I. Kenttämä, M. M. Abu-Omar, *ChemSusChem* **2014**, *7*, 2742–2747; d) D. Liu, E. Y.-X. Chen, *ACS Catal.* **2014**, *4*, 1302–1310.
- [18] a) T. Lu, X. Li, L. Gu, Y. Zhang, *ChemSusChem* **2014**, *7*, 2423–2426; b) L. Gu, T. Lu, X. Li, Y. Zhang, *Chem. Commun.* **2014**, *50*, 12308–12310.
- [19] a) H. Stetter and M. Schreckenberger, *Angew. Chem., Int. Ed. Engl.*, **1973**, *12*, 81–81. b) H. Stetter, *Angew. Chem., Int. Ed. Engl.*, **1976**, *15*, 639–647. c) H. Stetter and H. Kuhlmann, *Angew. Chem.*, **1974**, *86*, 589–589.
- [20] a) J. R. de Alaniz, M. S. Kerr, J. L. Moore and T. Rovis, *J. Org. Chem.*, **2008**, *73*, 2033–2040. b) T. Soeta, Y. Tabatake and Y. Ukaji, *Tetrahedron*, **2012**, *68*, 10188–10193. c) Y. Bao, N. Kumagai and M. Shibasaki, *Tetrahedron: Asymmetry*, **2014**, *25*, 1401–1408.
- [21] a) D. Enders, J. Han and A. Henseler, *Chem. Commun.*, **2008**, 3989–3991. b) Q. Liu, S. Perreault and T. Rovis, *J. Am. Chem. Soc.*, **2008**, *130*, 14066–14067. c) N. E. Wurz, C. G. Daniluc and F. Glorius, *Chem. - Eur. J.*, **2012**, *18*, 16297–16301.
- [22] J. Zhang, C. Xing, B. Tiwari and Y. R. Chi, *J. Am. Chem. Soc.*, **2013**, *135*, 8113–8116.
- [23] K. M. Doll, G. B. Bantchev and R. E. Murray, *ACS Sustainable Chem. Eng.*, **2013**, *1*, 39–45.
- [24] N. D. Vu, B. Guicheret, N. Duguet, E. Metay, M. Lemaire, *Green Chem.* **2017**, *19*, 3390–3399.
- [25] a) E. Deruer, N. Duguet, M. Lemaire, *ChemSusChem* **2015**, *8*, 2481–2486; b) N. D. Vu, S. Bah, E. Deruer, N. Duguet, M. Lemaire, *Chem. Eur. J.*, **2018**, *24*, 8141–8150.
- [26] a) L. Novák, G. Baán, J. Marosfalvi and C. Szántay, *Tetrahedron Lett.*, **1978**, *19*, 487–490. b) L. Novák, G. Baán, J. Marosfalvi and C. Szántay, *Chem. Ber.*, **1980**, *113*, 2939–2949.
- [27] R. Breslow, *J. Am. Chem. Soc.*, **1958**, *80*, 3719–3726.
- [28] S. M. Langdon, C. Y. Legault and M. Gravel, *J. Org. Chem.*, **2015**, *80*, 3597–3610.
- [29] a) G.-Q. Li, L.-X. Dai, S.-L. You, *Chem. Commun.* **2007**, 852–854; b) D. Enders, J. Han, A. Henseler, *Chem. Commun.* **2008**, 3989–3991; c) X. Bugaut, F. Liu, F. Glorius, *J. Am. Chem. Soc.* **2011**, *133*, 8130–8133; d) Z. Kelemen, O. Hollóczki, J. Nagy, L. Nyulászi, *Org. Biomol. Chem.* **2011**, *9*, 5362–5364; e) M. Padmanaban, A. T. Biju, F. Glorius, *Org. Lett.* **2011**, *13*, 98–101.
- [30] a) A. Miyashita, Y. Suzuki, Y. Okumura, T. Higashino, *Chem. Pharm. Bull.* **1996**, *44*, 252–254; b) A. Miyashita, Y. Suzuki, Y. Okumura, K. Iwamoto, T. Higashino, *Chem. Pharm. Bull.* **1998**, *46*, 6–11; c) Y. Suzuki, Y. Takemura, K. Iwamoto, T. Higashino, A. Miyashita, *Chem. Pharm. Bull.* **1998**, *46*, 199–206.
- [31] O. Hollóczki, Z. Kelemen, L. Nyulászi, *J. Org. Chem.* **2012**, *77*, 6014–6022.
- [32] The optimal temperature was found to be 130°C. See Supporting Information for details.
- [33] L. R. Domingo, P. Pérez, R. Contreras, *Tetrahedron*, **2004**, *60*, 6585–6591.
- [34] D. S. Allgäuer and H. Mayr, *Eur. J. Org. Chem.*, **2014**, 2956–2963.
- [35] D. P. Curran and Q. Zhang, *Adv. Synth. Catal.*, **2003**, *345*, 329–332.
- [36] a) C. E. I. Knappke, J. M. Neudörfl and A. Jacobi von Wangelin, *Org. Biomol. Chem.*, **2010**, *8*, 1695; b) C. E. I. Knappke, A. J. Arduengo III, H. Jiao, J. M. Neudörfl and A. Jacobi von Wangelin, *Synthesis*, **2011**, 3784–3795.
- [37] a) D. Enders, K. Breuer, G. Raabe, J. Runsink, J. H. Teles, J.-P. Melder, K. Ebel, S. Brode, *Angew. Chem. Int. Ed.* **1995**, *34*, 1021–1023; b) D. Enders, K. Breuer, J. Runsink, J. H. Teles, *Liebigs Ann.* **1996**, 2019–2028; c) S. Matsuoka, Y. Ota, A. Washio, A. Katada, K. Ichioka, K. Takagi and M. Suzuki, *Org. Lett.*, **2011**, *13*, 3722–3725; d) A. T. Biju, M. Padmanaban, N. E. Wurz and F. Glorius, *Angew. Chem., Int. Ed.*, **2011**, *50*, 8412–8415; e) Y. Zhang and E. Y.-X. Chen, *Angew. Chem., Int. Ed.*, **2012**, *51*, 2465–2469; f) M. Hong and E. Y.-X. Chen, *Angew. Chem., Int. Ed.*, **2014**, *53*, 11900–11906.
- [38] A. Bhunia, S. Thorat, R. G. Gonnade, A. T. Biju, *Chem. Commun.*, **2015**, *51*, 13690–13693.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

Vegetable oil-derived methyl azelaaldehydate (methyl 9-oxo-nonanoate) was converted to monomers through NHC-catalysed Stetter reaction, with limitation of the competing Benzoin condensation.

Organocatalysis*

*Aubin Charvieux, Nam Duc Vu, Nicolas Duguet, * Marc Lemaire**

Page No. – Page No.

Valorization of methyl azelaaldehydate – a vegetable oil-based platform molecule – to monomers through Stetter reaction

*one or two words that highlight the emphasis of the paper or the field of the study

Layout 2:

COMMUNICATION

((Insert TOC Graphic here; max. width: 11.5 cm; max. height: 2.5 cm; NOTE: the final letter height should not be less than 2 mm.))

Text for Table of Contents

Organocatalysis*

*Aubin Charvieux, Nam Duc Vu, Nicolas Duguet, * Marc Lemaire**

Page No. – Page No.

Valorization of methyl azelaaldehydate – a vegetable oil-based platform molecule – to monomers through Stetter reaction

*one or two words that highlight the emphasis of the paper or the field of the study